

The Number of Spanning Trees in Apollonian Networks

Zhongzhi Zhang, Bin Wu

*School of Computer Science and Shanghai Key Lab of Intelligent Information Processing,
Fudan University, Shanghai 200433, China (zhangzz@fudan.edu.cn).*

Francesc Comellas

*Dep. Matemàtica Aplicada IV, EETAC, Universitat Politècnica de Catalunya, c/ Esteve
Terradas 5, Castelldefels (Barcelona), Catalonia, Spain (comellas@ma4.upc.edu).*

Abstract

In this paper we find an exact analytical expression for the number of spanning trees in Apollonian networks. This parameter can be related to significant topological and dynamic properties of the networks, including percolation, epidemic spreading, synchronization, and random walks. As Apollonian networks constitute an interesting family of maximal planar graphs which are simultaneously small-world, scale-free, Euclidean and space filling and highly clustered, the study of their spanning trees is of particular relevance. Our results allow also the calculation of the spanning tree entropy of Apollonian networks, which then we compare with those of other graphs with the same average degree.

Keywords: Apollonian networks, small-world graphs, complex networks, self-similar, maximally planar, scale-free

1. Introduction

In the process known as Apollonian packing [10], which dates back to Apollonius of Perga (c262–c190 BC), we start with three mutually tangent circles, and draw their inner Soddy circle (tangent to the three circles). Next we draw the inner Soddy circles of this circle with each pair of the original three, and the process is iterated, see Fig. 1. An Apollonian packing can be used to design a graph, when each circle is associated to a vertex of the graph and vertices are connected if their corresponding circles are tangent. This graph, known as Apollonian graph or two-dimensional Apollonian network,

was introduced by Andrade et al. [1] and independently proposed by Doye and Massen in [7].

Apollonian networks have received much attention in recent years as two-dimensional Apollonian networks are scale-free, small-world, maximally planar, Euclidean and space filling [1, 30]. Dynamical processes taking place on these networks, such as percolation, epidemic spreading, synchronization and random walks, have been also investigated, see [30, 28, 29, 11, 31]. Some authors even suggest that the topological and dynamical properties of Apollonian networks are characteristic of neuronal networks as in the brain cortex [17].

In this paper, we study the number of spanning trees of two-dimensional Apollonian networks. This study is relevant given the importance of the graphs, and because the number of spanning trees of a finite graph is a graph invariant which characterizes the reliability of a network [6] and is related to its optimal synchronization and the study of random walks [14]. The number of spanning trees of a graph can be obtained from the product of all nonzero eigenvalues of the Laplacian matrix of the graph [9] (Kirchhoff's matrix-tree theorem). However, although this result can be applied to any graph, this calculation is analytically and computationally demanding.

Here, we follow a similar approach than for the Sierpiński gasket and Hanoi graphs [5, 20, 21, 25] and our method to compute the number of spanning trees in Apollonian networks is based on the self-similarity of the graphs.

2. Apollonian networks

We recall in this section the definition and main topological properties of two dimensional Apollonian networks. We use standard graph terminology and the words “network” and “graph” indistinctly.

Definition 2.1. *An Apollonian network $A(n)$, $n \geq 0$, is a graph constructed as follows:*

For $t = 0$, $A(0)$ is the complete graph K_3 (also called a 3-clique or triangle).

For $n \geq 1$, $A(n)$ is obtained from $A(n - 1)$: For each of the existing subgraphs of $A(n - 1)$ that is isomorphic to a 3-clique and created at step $n - 1$, a new vertex is introduced and connected to all the vertices of this subgraph. Figure 2 shows this construction process.

We note that Apollonian networks belong to the family of graphs known as k -trees, introduced in the 1970s [4], and which have been extensively




Figure 1: First stages of an Apollonian packing process.




Figure 2: Apollonian graphs $A(n)$ produced at iterations $n = 0, 1, 2, 3$ and 4.

studied since. They are defined as follows: a k -clique is a k -tree; if G is a k -tree, adding a vertex to G and joining it to all vertices of an existing k -clique in G yields a new graph G' , which is also a k -tree. Thus, Apollonian networks are k -trees, since they start with a 3-clique (that is, the only possible k -tree with 3 vertices), and since each time a vertex is added, it is joined to all the vertices of an existing 3-clique. k -trees are of interest as they are maximal graphs for a given tree-width.

Thanks to the deterministic nature of the graphs $A(n)$, it is possible to give exact values for relevant topological properties of these graphs, including degree distribution, degree correlations, clustering, diameter and average distance.

Order and size of $A(n)$.– The order and size of an Apollonian graph $A(n)$ are $V_n = |V(n)| = \frac{1}{2}(3^n + 5)$ and $E_n = |E(n)| = \frac{3}{2}(3^n + 1)$.

Planarity– A graph is planar if it can be drawn on the plane with no edges crossing. A planar graph is maximal, or maximally planar, if it cannot be extended to a larger planar graph by adding an edge.

Apollonian networks are maximally planar [1] and this is an important feature for its relevance in the development of efficient algorithms [3].

Degree distribution.– The cumulative degree distribution of an Apollonian network $A(n)$ is $P(k) \propto k^{1-\gamma}$. with $\gamma = \ln 3 / \ln 2 = 2.58496$, see [1, 7]. The graph is scale-free. Many real networks share this property with exponent values in the same range as $A(n)$ [15].

Correlation coefficient.– From the Pearson correlation coefficient for the degrees of the endvertices of the edges of $A(n)$, Doye and Massen obtained in [7] the exact value of the correlation coefficient and it is always negative and thus the network is disassortative. For large values of n , $r(n) \sim -\frac{10}{3} \left(\frac{2}{3}\right)^n$. Thus, this parameter goes to zero as the order of the graph increases. Technological and biological networks are usually disassortative as it is also the case of some information networks, see [15, 19].

Modularity and self-similarity.– From Fig. 2, we see that Apollonian networks are self-similar, suggesting an alternative way to construct them. As shown in Fig. 3, $A(n+1)$ can be obtained by joining three replicas of $A(n)$, labeled by A_n^1 , A_n^2 and A_n^3 , and identifying three pairs of edges. Thus, the graphs are modular. Modularity can be quantified with the function Q introduced by Newman and Girvan [16]. For Apollonian networks $Q = 0.5938$. Moreover, the communities are spatially localized [1, 7].




Figure 3: Recursive construction of Apollonian networks, pointing out their self-similarity. $A(n+1)$ can be obtained by joining three replicas of $A(n)$, labeled here A_n^1 , A_n^2 and A_n^3 , after merging three pairs of edges.

Clustering.— The clustering coefficient C of $A(n)$ is 0.8284 for large n . Furthermore, the clustering coefficient of a vertex of $A(n)$, for large n , is inversely proportional to its degree [1, 28]: $C(k) = \frac{4(k - \frac{3}{2})}{k(k - 1)}$. This property is considered to be a signature of a hierarchical network structure, see [2]

Average distance and diameter.— The modular recursive construction of $A(n)$ allows to obtain the exact analytical value for its average distance [24]. The average distance of $A(n)$, for n large, follows $\bar{d}(n) \sim \ln |V_n|$ which shows a logarithmic scaling with the order of the graph. As the diameter has a similar behavior [23], the graph is small-world.

3. The number of spanning trees in Apollonian networks

In this section, we find the number of spanning trees of the Apollonian network $A(n)$. For this calculation we use a decimation method [12] which has been used to find the number of spanning trees in other recursive graph families like the Sierpiński gasket [5], the pseudofractal web [26], and some fractal lattices [27]. The particular structure of Apollonian networks, allow us to write recursive equations for the number of spanning trees which are solved by induction.

In the following, we denote by V_n and E_n the number of vertices and edges of $A(n)$. Then a spanning subgraph of $A(n)$ is a subgraph with the same vertex set as $A(n)$ and a number of edges E'_n such that $E'_n \leq E_n$. A spanning tree of $A(n)$ is a spanning subgraph which is a tree and thus $E'_n = V_n - 1$.

We call “hub vertices” the three outmost vertices in the construction as

shown in Fig. 2 and “hub edges” the three exterior edges which connect the hub vertices.

To simplify our calculations, we define some quantities as follows: We denote as s_n the total number of spanning trees of $A(n)$. a_n represents the number of spanning subgraphs of $A(n)$ which consist of three trees such that each hub vertex of $A(n)$ belongs to a different tree. In a similar way, b_n will denote the number of spanning subgraphs of $A(n)$ which consist of two trees such that no hub edges belong to the spanning subgraph and one of the hub vertices of the subgraph belongs to one tree and the other two hub vertices are in the second tree. In a similar way, and taking into account the tree to which a given hub vertex belongs, we define b'_n and b''_n . Finally, c_n is the number of spanning trees of $A(n)$ which have no hub edges. We notice that the equation $b_n = b'_n = b''_n$ holds as a result of symmetry, thus, in what follows, we will use b_n instead of b'_n and b''_n . Figure 4 illustrates schematically these five types of spanning subgraphs which we denote in all figures as A_n , B_n , B'_n , B''_n , C_n , respectively.




Figure 4: Schematic illustration of the five types of spanning subgraphs A_n , B_n , B'_n , B''_n and C_n derived from $A(n)$. In this figure, vertices represent the hub vertices of $A(n)$ and two hub vertices joined by a black line belong to different trees. A red line means that the two hub vertices are in the same tree but have no hub edge joining them.

In the previous definitions, we have not considered the cases where the spanning subgraph contains hub edges. We deal with these cases in the following three lemmas.

Lemma 3.1. *The number of spanning subgraphs of $A(n)$ which consist of two trees such that one hub edge with its two hub vertices belongs to one tree while the third hub vertex of $A(n)$ is in the other tree equals a_n .*

Proof. Let $\bar{B}_n$ be the set of subgraphs considered in this Lemma. We verify the correctness of the result by showing that there exists an one-to-one correspondence between the set $\bar{B}_n$ and A_n as it is shown in Fig. 5. For every spanning subgraph in $\bar{B}_n$, if we remove the hub edge, then the three hub vertices will belong to three different trees, so it belongs to A_n . Conversely, for every spanning subgraph in A_n , if we add a hub edge, then its




Figure 5: Graphical illustration of Lemma 3.1. The green line means that the hub edge joining these two hub vertices belongs to the spanning subgraph.

two hub vertices belong to one tree and the subgraph is in $\bar{B}_n$. Thus, there exists a one-to-one correspondence between $\bar{B}_n$ and A_n , and the cardinality of $\bar{B}_n$ is a_n . $\square$

Lemma 3.2. *The number of spanning subgraphs of $A(n)$ such that they contain just one hub edge and one hub vertex which is connected to one of the hub vertices of this edge through edges of the tree is b_n .*

Proof. Let $\bar{C}_n$ be the set of subgraphs considered in this Lemma. As



Figure 6: Graphical illustration of Lemma 3.2. The green line means that the hub edge joining these two hub vertices belongs to the spanning subgraph.

before, we prove this Lemma by showing that sets $\bar{C}_n$ and B_n'' have an one-to-one correspondence. This can be verified by deleting the hub edge, see Fig. 6. Thus, the cardinality of $\bar{C}_n$ is b_n'' . $\square$

Lemma 3.3. *The number of spanning subgraphs of $A(n)$ that include two hub edges is a_n .*

Proof. Consider the bijection between $\bar{C}_n'$, the set of subgraphs which contain two hub edges, and A_n (Fig. 7). $\square$

Next we establish a recursive relationship among the parameters s_n , a_n , b_n , and c_n .




Figure 7: Graphical illustration of Lemma 3.3. The green line means that the hub edge joining these two hub vertices belongs to the spanning subgraph .

Lemma 3.4. For $n \geq 0$, $a_{n+1} = 3a_n^3 + 6a_n^2b_n$.

Proof. We prove this result by considering a graphical version of the equation (Fig. 8) which represents the recursive construction method of $A(n+1)$ from $A(n)$ and enumerates all possible contributions to a_{n+1} .

In this representation we only draw four vertices in each case, since each non drawn (interior) vertex connects at least to one of these four vertices (although they do not have necessarily to be adjacent). This is sufficient to determine whether each case belongs to A_{n+1} , B_{n+1} or C_{n+1} .




Figure 8: Configurations needed to find a_{n+1} from a_n and b_n . In this representation, the red curve denotes the spanning tree to which the two hub vertices belong to. The number on the right of the figure counts configurations that, by symmetry, contribute to a_{n+1} through the merging process

Next we should prove that each configuration is correct, but we only analyze in detail the first additive term as the other term can be verified in a similar way. For this case (see Fig. 9), hub vertices h_1 and h_4 are connected while h_2 and h_3 are not. Thus, after merging, there are three spanning trees and we confirm that this subgraph belongs to A_{n+1} . Because of the symmetry, h_4 can also be connected to h_2 or h_3 and we count three times this case. $\square$

Lemma 3.5. For $n \geq 0$, $b_{n+1} = a_n^3 + 7a_n^2b_n + 7a_nb_n^2 + a_n^2c_n$.

Proof. We prove the lemma by enumeration. Figure 10 shows all the




Figure 9: The first configuration which contributes to a_{n+1} .




Figure 10: Configurations needed to find b_{n+1} .

distinct possibilities. Again, we only analyze the first case. We label the four hub vertices in the same way as in Fig. 9. In the first case, h_1 , h_2 , h_4 are all connected while h_3 is not. There are two spanning trees, and one has no hub edges, so this configuration belongs to set B_{n+1} . Symmetries generate equivalent configurations and the factor is one. $\square$

Lemma 3.6. For $n \geq 0$, $c_{n+1} = a_n^3 + 12a_n^2b_n + 36a_nb_n^2 + 14b_n^3 + 3a_n^2c_n + 12a_nb_nc_n$.

Proof. As in former lemmas, the proof is by enumeration of all possible contributions to c_{n+1} , see in Fig. 11 the details. In the first case, h_1 , h_2 , h_3 and h_4 are all connected and the merging process produces a spanning tree. As no hub edges are included in it, we can see that this case belongs to set C_{n+1} . Besides, because of the symmetry, only this configuration is relevant. All other cases are analyzed similarly and we omit the details. $\square$

Lemma 3.7. For $n \geq 0$, $s_{n+1} = 16a_n^3 + 72a_n^2b_n + 78a_nb_n^2 + 14b_n^3 + 9a_n^2c_n + 12a_nb_nc_n$.

Proof. Figure 12 shows all the configurations contributing s_{n+1} . We do not give the calculation details as they are like in former lemmas. $\square$




Figure 11: Configurations that contribute to c_{n+1} .




Figure 12: Illustration for the calculation of s_{n+1} . Here, a blue line indicates that the corresponding hub vertices belong to the same spanning tree (including both the cases where they are adjacent and they are connected through other vertices).

Lemma 3.8. For $n \geq 0$, $a_n c_n = 3b_n^2$.

Proof. By induction. For $n = 0$, using the initial conditions $a_0 = 1$, $b_0 = 0$ and $c_0 = 0$, the result is true.

Then we assume that for $n = k$, the lemma is true. For $n = k + 1$, and using Lemmas 3.4-3.6, we have that

$$\begin{aligned} a_{k+1}c_{k+1} - 3b_{k+1}^2 &= (3a_k^3 + 6a_k^2b_k)(a_k^3 + 12a_k^2b_k + 36a_kb_k^2 + 14b_k^3 + 3a_k^2c_k \\ &\quad + 12a_kb_kc_k) - 3(a_k^3 + 7a_k^2b_k + 7a_kb_k^2 + a_k^2c_k)^2 \\ &= 3a_k^2(a_k^2 + 4a_kb_k + 7b_k^2 - a_kc_k)(a_kc_k - 3b_k^2), \end{aligned}$$

As by induction hypothesis, we have that $a_kc_k - 3b_k^2 = 0$ we obtain the result. $\square$

Lemma 3.9. For $n \geq 0$, $\frac{a_{n+1}}{a_n^3} = \frac{5^n}{3^{n-1}}$.

Proof. Define $f_n = \frac{a_{n+1}}{a_n^3}$. From Lemma 3.4, we have

$$\frac{a_{n+1}}{a_n^3} = \frac{3a_n^3 + 6a_n^2b_n}{a_n^3} = 3 + 6\frac{b_n}{a_n},$$

and thus

$$\frac{b_n}{a_n} = \frac{1}{6} \left(\frac{a_{n+1}}{a_n^3} - 3 \right).$$

We use Lemmas 3.8 and 3.5, to obtain:

$$\begin{aligned} \frac{b_{n+1}}{a_n^3} &= 1 + 7\frac{b_n}{a_n} + 10 \left(\frac{b_n}{a_n} \right)^2 \\ &= 1 + 7 \left[\frac{1}{6} \left(\frac{a_{n+1}}{a_n^3} - 3 \right) \right] + 10 \left[\frac{1}{6} \left(\frac{a_{n+1}}{a_n^3} - 3 \right) \right]^2 \\ &= -\frac{a_{n+1}}{2a_n^3} + \frac{5a_{n+1}^2}{18a_n^6}, \end{aligned}$$

which gives:

$$\frac{b_{n+1}}{a_{n+1}} = \frac{b_{n+1}}{a_n^3} \frac{a_n^3}{a_{n+1}} = \left(-\frac{a_{n+1}}{2a_n^3} + \frac{5a_{n+1}^2}{18a_n^6} \right) \frac{a_n^3}{a_{n+1}} = -\frac{1}{2} + \frac{5a_{n+1}}{18a_n^3}.$$

Thus

$$\frac{a_{n+2}}{a_{n+1}^3} = 3 + 6\frac{b_{n+1}}{a_{n+1}} = 3 + 6 \left(-\frac{1}{2} + \frac{5a_{n+1}}{18a_n^3} \right) = \frac{5a_{n+1}}{3a_n^3},$$

which we can write as

$$f_{n+1} = \frac{5}{3}f_n.$$

From the initial condition $f_0 = \frac{a_1}{a_0^3} = 3 = \frac{5^0}{3^{-1}}$, we finally find

$$f_n = \frac{a_{n+1}}{a_n^3} = \frac{5^n}{3^{n-1}}.$$

□

Lemma 3.10. For $n \geq 0$, $a_n = 3^{-\frac{1}{4} + \frac{3^n}{4} + \frac{n}{2}} 5^{-\frac{1}{4} + \frac{3^n}{4} - \frac{n}{2}}$.

Proof. From the former Lemma we have $a_{n+1} = \frac{5^n}{3^{n-1}} a_n^3$, which with initial condition $a_0 = 1$ gives

$$a_n = 3^{-\frac{1}{4} + \frac{3^n}{4} + \frac{n}{2}} 5^{-\frac{1}{4} + \frac{3^n}{4} - \frac{n}{2}}.$$

□

Lemma 3.11. For $n \geq 0$, $b_n = \frac{1}{2} 15^{\frac{1}{4}(-1+3^n-2n)} (5^n - 3^n)$.

Proof. From Lemma 3.4 we know that $b_n = \frac{a_{n+1} - 3a_n^3}{6a_n^2}$ and using Lemma 3.10 we obtain

$$b_n = \frac{1}{2} 15^{\frac{1}{4}(-1+3^n-2n)} (5^n - 3^n).$$

□

Lemma 3.12. For $n \geq 0$, $c_n = \frac{1}{4} 3^{\frac{1}{4}(3+3^n-6n)} 5^{\frac{1}{4}(-1+3^n-2n)} (3^n - 5^n)^2$.

Proof. According to Lemma 3.8, we have $c_n = \frac{3b_n^2}{a_n}$, and using the results of Lemmas 3.10 and 3.11, we obtain

$$c_n = \frac{1}{4} 3^{\frac{1}{4}(3+3^n-6n)} 5^{\frac{1}{4}(-1+3^n-2n)} (3^n - 5^n)^2.$$

□

Theorem 3.13. The number of spanning trees of $A(n)$ is

$$s_n = \frac{1}{4} 3^{\frac{3}{4}(-1+3^{n-1}-2(n-1))} 5^{\frac{1}{4}(-3+3^n-2(n-1))} (3^n + 5^n)^2.$$

Proof. The result comes from Lemmas 3.7, 3.10, 3.11 and 3.12. $\square$

After having an explicit expression for the number of spanning trees of $A(n)$, we can calculate its spanning tree entropy, which is defined as in [22, 13]:

$$z = \lim_{n \rightarrow \infty} \frac{\ln s_n}{V_n}.$$

The spanning tree entropy is an interesting parameter characterizing the network structure.

Corollary 3.14. *The spanning tree entropy of Apollonian networks is $\frac{\ln 15}{2}$.*

Proof. Define $z_n = \frac{\ln s_n}{V_n}$. From Theorem 3.13, we have

$$z_n = \frac{-8 \ln 2 + \ln \frac{27}{5} + 3^n \ln 15 - 2n \ln 135 + 8 \ln(3^n + 5^n)}{2(5 + 3^n)},$$

and finally

$$z = \lim_{n \rightarrow \infty} z_n = \frac{\ln 15}{2}.$$

$\square$

We can compare this asymptotic value of the entropy of the spanning trees for Apollonian networks, $z = \frac{\ln 15}{2} \simeq 1.3540$, with that of other graphs with the same average degree. For example, the value for the Sierpinski graph is 1.5694 [5] and for the 3-dimensional hypercubic lattice $\mathcal{L}_3$ is 1.6734 [18, 8]. Thus, the asymptotic value for Apollonian networks is the lowest reported for graphs with average degree 6. This reflects the fact that the number of spanning trees in $A(n)$, although growing exponentially, do it at a lower rate than graphs with the same average degree.

4. Conclusion

In this paper we find the number of spanning trees in Apollonian networks by using a method, based on its self-similar structure, which allows us to obtain an exact analytical expression for any number of discs. The method could be used to further study in this graph, and other self-similar graphs, their spanning forests, connected spanning subgraphs, random walks and vertex or edges coverings. Knowing the number of spanning trees for Apollonian networks allows us to show that their spanning tree entropy is lower than in other graphs with the same average degree.

Acknowledgements

Z. Zhang is supported by the National Natural Science Foundation of China under Grants No. 61074119. F. Comellas is supported by the Ministerio de Economía y Competitividad, Spain, and the European Regional Development Fund under project MTM2011-28800-C02-01 and partially supported by the Catalan Research Council under grant 2009SGR1387.

References

- [1] J. S. Andrade Jr., H. J. Herrmann, R. F. S. Andrade, L. R. da Silva, Apollonian Networks: Simultaneously scale-free, small world, Euclidean, space filling, and with matching graphs, *Phys. Rev. Lett.* 94 (2005) 018702.
- [2] A. L. Barabási, Z.N. Oltvai, Network biology: Understanding the cell's functional organization, *Nat. Rev. Genet.* 5 (2004) 101-114.
- [3] A. Brandstädt, V. B. Le, J. P. Spinrad. Graph Classes: A Survey (SIAM Monographs on Discrete Mathematics and Applications) Philadelphia, PA, 1999.
- [4] L.W. Beineke, R.E. Pippert, Properties and characterization of k -trees, *Mathematika* 18 (1971) 141–151.
- [5] S.-C. Chang, L.-C. Chen, W.-S. Yang, Spanning trees on the Sierpiński gasket, *J. Stat. Phys.* 126 (2007) 649–667.
- [6] C.J. Colbourn, The Combinatorics of Network Reliability, New York, Oxford University Press 1987.
- [7] J. P. K. Doye, C. P. Massen, Self-similar disk packings as model spatial scale-free networks, *Phys. Rev. E* 71 (2005) 016128.
- [8] J.H. Felker, R. Lyons, High-precision entropy values for spanning trees in lattices, *J. Phys. A: Math. Gen.* 36 (2003) 83618365.
- [9] C. Godsil, G. Royle, Algebraic Graph Theory, Graduate Texts in Mathematics 207, Springer, New York 2001.
- [10] E. Kasner, F.D Supnick, The Apollonian packing of circles, *Proc. Natl. Acad. Sci. U.S.A.* 29 (1943) 378–384.

- [11] Z-G. Huang, X-J. Xu, Z-X. Wu, Y.H. Wang, Walks on Apollonian networks, *Eur. Phys. J. B* 51 (2006) 549–553.
- [12] M. Knezevic, J. Vannimenus, Large-scale properties and collapse transition of branched polymers: Exact results on fractal lattices, *Phys. Rev. Lett.* 56 (1986) 1591.
- [13] R. Lyons, Asymptotic enumeration of spanning trees, *Combin. Probab. Comput.* 14 (2005) 491–522.
- [14] P. Marchal, Loop-erased random walks, spanning trees and hamiltonian cycles, *Elect. Comm. in Probab.* 5 (2000) 39-50.
- [15] M.E.J. Newman, The structure and function of complex networks, *SIAM Review* 45 (2003) 167–256.
- [16] M. E. J. Newman and M. Girvan, Finding and evaluating community structure in networks, *Phys. Rev. E* 69 (2004) :026113.
- [17] G.L. Pellegrini, L. de Arcangelis, H.J. Herrmann, C. Perrone-Capano. Modelling the brain as an Apollonian network, arXiv:q-bio/0701045v1, 2007.
- [18] R. Shrock and F. Y. Wu, Spanning trees on graphs and lattices in d dimensions, *J. Phys. A: Math. Gen.* 33 (2000) 3881.
- [19] R. V. Solé, S. Valverde, Information theory of complex networks: on evolution and architectural constraints, *Lecture Notes in Phys.* 650 (2004) 189–207.
- [20] E. Teufl, S. Wagner, Resistance scaling and the number of spanning trees in self-similar lattices, *J. Stat. Phys.* 142 (2011) 879–897.
- [21] E. Teufl, S. Wagner, The Number of Spanning Trees in Self-Similar Graphs, *Ann. Comb.* 15 (2011) 355–380.
- [22] F. Y. Wu, Number of spanning trees on a lattice, *J. Phys. A: Math. Gen.* 10 (1977) 113-115.
- [23] Z. Zhang, F. Comellas, G. Fertin, L. Rong, High dimensional Apollonian networks, *J. Phys. A* 39 (2006) 1811–1818.
- [24] Z. Zhang, L. Chen, S. Zhou, L. Fang, J. Guan, T. Zou, Analytical solution of average path length for Apollonian networks, *Phys. Rev. E* 77 (2008) 017102.

- [25] Z. Zhang, S. Wu, F. Comellas, The number of spanning trees of Hanoi graphs, submitted.
- [26] Z. Zhang, H. Liu, B. Wu, S. Zhou, Enumeration of spanning trees in a pseudofractal scale-free web, *Europhys. Lett.* 90 (2010) 68002.
- [27] Z. Zhang, H. Liu, B. Wu, T. Zou, Spanning trees in a fractal scale-free lattice, *Phys. Rev. E* 83 (2011) 016116.
- [28] Z. Zhang, L. Rong, F. Comellas, High dimensional random Apollonian networks, *Physica A* 364 (2005) 610–618.
- [29] Z. Zhang, L. Rong, S.G. Zhou, Evolving Apollonian networks with small-world scale-free topologies, *Phys. Rev. E* 74 (2006) 046105.
- [30] T. Zhou, G. Yan, and B.H. Wang, Maximal planar networks with large clustering coefficient and power-law degree distribution, *Phys. Rev. E* 71 (2005) 046141.
- [31] Z. Zhang, S. Zhou, Correlations in random Apollonian networks, *Physica A* 380 (2007) 621–628.